



ELSEVIER

Topology and its Applications 69 (1996) 153–164

---

---

**TOPOLOGY  
AND ITS  
APPLICATIONS**

---

---

## Solutions to two problems of J.D. Lawson and M. Mislove<sup>☆</sup>

Liu Ying-Ming, Liang Ji-Hua<sup>\*</sup>

*Institute of Mathematics, Sichuan University, Chengdu, China*

Received 4 January 1995; revised 21 March 1995, 26 July 1995

---

### Abstract

The problem of characterizing those  $L$  for which  $[X \rightarrow L]$  is a continuous DCPO for all core compact spaces is solved. Also, the problem of coincidence of the Isbell topology with the Scott topology on  $[X \rightarrow L]$  is solved.

**Keywords:** Continuous DCPO; L-Domain; Core compact space; Isbell topology

**AMS classification:** 54A10; 06B35

---

### 1. Introduction and preliminaries

Let  $X$  be a topological space and  $L$  a DCPO (i.e.,  $L$  is a partially ordered set in which every directed set has the supremum) equipped with the Scott topology  $\sigma(L)$ . Write  $[X \rightarrow L]$  for the set of all continuous mappings from  $X$  to  $L$ ; then the set  $[X \rightarrow L]$  with the pointwise order is again a DCPO. Recently, in [1] Lawson and Mislove posed the following problems:

**Problem A.** Characterize those  $L$  for which  $[X \rightarrow L]$  is a continuous DCPO for all core compact spaces  $X$ . A likely candidate is the class of all continuous L-Domains  $L$ . Does one get the same answer if one restricts to the core compact spaces which are also compact?

**Problem B.** Let  $X$  be a core compact space and  $L$  be a DCPO equipped with the Scott topology. Under what conditions on  $L$  do the Isbell and Scott topologies on  $[X \rightarrow L]$  agree?

---

<sup>☆</sup> The project is supported by National Science Foundation of China.

<sup>\*</sup> Corresponding author.

The aim of this paper is to give a complete solution to Problem A and a partial solution to Problem B. Firstly, we point out that the last question in Problem A has a negative answer and hence Problem A is solved by means of the following theorem.

**Theorem A.** *For a DCPO  $L$ , the following conclusions are equivalent:*

- (1)  $L$  is a continuous  $L$ -Domain;
- (2)  $[X \rightarrow L]$  is a continuous  $L$ -Domain for all core compact spaces  $X$ ;
- (3)  $[X \rightarrow L]$  is a continuous DCPO for all compact spaces  $X$ .

Then two examples are given to show that Problem B has no positive answer even for continuous  $L$ -Domains with least element and bounded complete continuous DCPOs without least element. Finally, the following conclusion is proved and hence Problem B is solved.

**Theorem B.** *Let  $L$  be a continuous  $L$ -Domain with least element  $O_L$ . Then the Isbell and Scott topologies on  $[X \rightarrow L]$  agree for all core compact spaces  $X$  if and only if  $L$  is a bounded complete DCPO.*

For a DCPO  $L$  and  $x, y \in L$ , we say that  $x$  is way below  $y$  ( $x \ll y$ ), if given a directed set  $D \subseteq L$  with  $y \leq \bigvee D$  then  $x \leq d$  for some  $d \in D$ . A DCPO  $L$  is said to be continuous, if for all  $x \in L$  the set  $\downarrow x = \{y \in L: y \ll x\}$  is directed and  $x = \bigvee \downarrow x$ . If  $L$  is simultaneously a complete lattice and a continuous DCPO, then  $L$  is called a continuous lattice.

The following facts are well-known and often used in the present paper.

**Theorem 1.1** [2]. *In any continuous DCPO  $L$ , we have the following conclusions:*

- (1)  $x \ll y$  implies there is a  $z \in L$  such that  $x \ll z \ll y$ ;
- (2)  $\{\uparrow x = \{y \in L: y \gg x\}: x \in L\}$  is a base of the Scott topology  $\sigma(L)$ ;
- (3) the Scott topology  $\sigma(L)$  is a continuous lattice.

A DCPO  $L$  is a  $L$ -Domain if for all  $x \in L$  the set  $\downarrow x = \{y \in L: y \leq x\}$  is a complete lattice;  $L$  is called bounded complete if  $L$  is a DCPO and each nonempty subset with upper bound has a supremum. Obviously, a bounded complete DCPO is also a  $L$ -Domain if  $\downarrow x$  has a least element for each  $x \in L$ , but the reverse is not true.

We say that a topological  $X$  is core compact if the lattice  $\Omega(X)$  of its open sets is a continuous lattice.

## 2. Solution to Problem A

In the following, for a DCPO  $L$  and  $x \in L$  we write  $\bigvee_x$  for the supremum operation in  $\downarrow x$  and  $O_x$  for the least element of  $\downarrow x$  if it exists, and  $\mathbb{N}$  stands for the set of natural numbers.

**Lemma 2.1.** *In any L-Domain  $L$ , one has the following statements for each  $x \in L$ :*

- (1)  $O_x$  is a minimal element of  $L$  and hence  $O_x \ll O_x$ ;
- (2) if  $\{a_i: i \in \Delta\} \subseteq L$  with  $\bigvee_{i \in \Delta} a_i \geq O_x$ , then  $O_{a_i} = O_x$  for all  $i \in \Delta$ .

**Proof.** It directly follows from the definition of  $O_x$ .  $\square$

For a core compact space  $X$  and a continuous DCPO  $L$  with least element  $O_L$ , we can easily show, as in [1],  $\bigvee \downarrow f = f$  for each  $f \in [X \rightarrow L]$  by constructing mappings that take some constant value on an open set of  $X$  and value  $O_L$  otherwise. For a general continuous DCPO, it seems more difficult to show this. The following construction is essential.

For  $A \subseteq X$ ,  $a \in L$  and  $f \in [X \rightarrow L]$ , define a mapping

$$F(a, A, f): X \rightarrow L \quad \text{by } F(a, A, f)(x) = \begin{cases} a, & x \in A, \\ O_{f(x)}, & x \notin A, \end{cases}$$

and write  $f_{\langle\langle a \rangle\rangle} = f^{-1}(\uparrow a)$  for short.

**Lemma 2.2.** *Let  $L$  be a continuous L-Domain,  $a \in L$  and  $f \in [X \rightarrow L]$ . Then*

$$F(a, U, f) \in [X \rightarrow L]$$

*if  $U \in \Omega(X)$  and  $U \subseteq f_{\langle\langle a \rangle\rangle}$ .*

**Proof.** Take an arbitrary  $b \in L$  and note that  $f_{\langle\langle a \rangle\rangle} \cap f_{\langle\langle b \rangle\rangle} = \emptyset$  if  $O_b \neq O_a$ . Then  $f_{\langle\langle b \rangle\rangle} \subseteq X \setminus U$  as  $U \subseteq f_{\langle\langle a \rangle\rangle}$ . By Lemma 2.1 we can verify that the following

$$F(a, U, f)_{\langle\langle b \rangle\rangle} = \begin{cases} f_{\langle\langle b \rangle\rangle} \cup U, & b = O_b \text{ and } a \gg b, \\ f_{\langle\langle b \rangle\rangle} \setminus U = f_{\langle\langle b \rangle\rangle}, & b = O_b \text{ and } a \notin \uparrow b, \\ U, & b \neq O_b \text{ and } a \gg b, \\ \emptyset, & b \neq O_b \text{ and } a \notin \uparrow b, \end{cases}$$

hold, i.e.,  $F(a, U, f)_{\langle\langle b \rangle\rangle} \in \Omega(X)$ . By Theorem 1.1  $F(a, U, f) \in [X \rightarrow L]$ .  $\square$

**Remark 2.3.** In this lemma, the condition  $U \subseteq f_{\langle\langle a \rangle\rangle}$  is necessary. For example, let  $L$  be the continuous L-Domain denoted by Fig. 1, and  $X = L$  equipped with the Scott topology and  $f$  the identity mapping on  $L$ . Take  $U = \{b_1, b_2, b_3\} \in \Omega(X)$ , then

$$F(a_3, U, f)(x) = \begin{cases} a_3, & x \in \{b_1, b_2, b_3\}, \\ O_a, & x \in \{a_1, a_2, a_3, O_a\}, \\ O_b, & x = O_b, \end{cases}$$

and hence  $F(a_3, U, f)_{\langle\langle b \rangle\rangle} = \{O_b\} \notin \Omega(X)$  for  $b = O_b$ , i.e.,  $F(a_3, U, f) \notin [X \rightarrow L]$ .

**Lemma 2.4.** *Let  $L$  be a continuous L-Domain,  $a, b \in L$  and  $V \in \Omega(X)$ ,  $f \in [X \rightarrow L]$ . Then  $F(b, V, f) \ll F(a, f_{\langle\langle a \rangle\rangle}, f)$  in  $[X \rightarrow L]$  if  $b \ll a$  and  $V \ll f_{\langle\langle a \rangle\rangle}$ .*




Fig. 1.

**Proof.** By Lemma 2.2, we have  $F(b, V, f), F(a, f_{\langle\langle a \rangle\rangle}, f) \in [X \rightarrow L]$ .

Now suppose that  $\{h_j: j \in \Delta\}$  is a directed family in  $[X \rightarrow L]$  with  $\bigvee_{j \in \Delta} h_j \geq F(a, f_{\langle\langle a \rangle\rangle}, f)$ ; then

$$\left(\bigvee h_j\right)_{\langle\langle b \rangle\rangle} = \bigcup_{j \in \Delta} h_{j \langle\langle b \rangle\rangle} \supseteq F(a, f_{\langle\langle a \rangle\rangle}, f)_{\langle\langle b \rangle\rangle} \supseteq f_{\langle\langle a \rangle\rangle}$$

by the proof of Lemma 2.2. Hence we have  $h_{j_0 \langle\langle b \rangle\rangle} \supseteq V$  for some  $j_0 \in \Delta$  because  $\{h_{j \langle\langle b \rangle\rangle}: j \in \Delta\}$  is directed. By Lemma 2.1(2)  $O_{f(x)} \leq h_{j_0}(x)$  for all  $x \in X \setminus V$ , hence  $F(b, V, f) \leq h_{j_0}$ . Thus the lemma is proved.  $\square$

**Lemma 2.5.** Let  $L$  be an  $L$ -Domain, then we have the following conclusions:

- (1)  $L$  is continuous if and only if  $\downarrow x$  is a continuous lattice for all  $x \in L$ .
- (2) If  $a, b \in \downarrow x \cap \downarrow y$  and  $y \geq a \bigvee_x b = a \bigvee_y b$ .

**Proof.** (1) For  $x \in L$  and  $a \in \downarrow x$ ,  $\downarrow a \subseteq \downarrow_x a = \{b \in \downarrow x: b \ll a \text{ in } \downarrow x\}$  is obvious and  $\downarrow_x a$  is directed because  $\downarrow x$  is a complete lattice, which implies that  $\downarrow x$  is a continuous lattice if  $L$  is continuous. Conversely we need only show that  $y \ll x$  in  $\downarrow x$  implies  $y \ll x$  in  $L$  for all  $x \in L$  and  $y \in \downarrow x$ . Suppose that  $y \ll x$  in  $\downarrow x$  and  $D$  is a directed set in  $L$  with  $\bigvee D \geq x$  and let  $e = \bigvee D$ . Then  $\downarrow_e x = \{b \in \downarrow e: b \ll x \text{ in } \downarrow e\} \subseteq \downarrow x$  is directed and  $\bigvee \downarrow_e x = x$  as  $\downarrow e$  is continuous, thus we have  $y \leq b$  for some  $b \in \downarrow_e x$ , i.e.,  $y \ll x$  in  $\downarrow e$ . As  $D$  is also a directed set in  $\downarrow e$  and  $\bigvee D = e \geq x$ , then  $y \leq d$  for some  $d \in D$ , which implies  $y \ll x$  in  $L$ .

(2)  $a \bigvee_x b$  is obviously an upper bound of  $\{a, b\}$  in  $\downarrow y$  by  $y \geq a \bigvee_x b$ , hence  $a \bigvee_x b \geq a \bigvee_y b$  holds. In this case we have  $a \bigvee_y b \leq x$ , which implies that  $a \bigvee_y b$  is also an upper bound of  $\{a, b\}$  in  $\downarrow x$ , so  $a \bigvee_y b \geq a \bigvee_x b$  holds and hence  $a \bigvee_x b = a \bigvee_y b$ .  $\square$

**Lemma 2.6.** Let  $L$  be a continuous  $L$ -Domain and  $X$  a core compact space. Then  $f = \bigvee \downarrow f$  for all  $f \in [X \rightarrow L]$ .

**Proof.** Note  $F(a, f_{\langle\langle a \rangle\rangle}, f) \leq f$  for all  $a \in L$ ; it is easy to show

$$f = \bigvee_{a \in L} \bigvee_{b \in \downarrow a} \bigvee_{V \ll f_{\langle\langle a \rangle\rangle}} F(b, V, f)$$

because both  $L$  and  $\Omega(X)$  are continuous. By Lemma 2.4 we obtain  $f = \bigvee \downarrow f$ .

**Lemma 2.7.**  $[X \rightarrow L]$  is an  $L$ -Domain for each continuous  $L$ -Domain  $L$  and core compact space  $X$ .

**Proof.** Take an arbitrary  $f \in [X \rightarrow L]$  and note that  $\downarrow f$  is a DCPO; it is sufficient to show that  $\downarrow f$  is a  $\vee$ -semilattice with least element. First of all we define the mapping  $O_f: X \rightarrow L$  by  $O_{f(x)} = O_f(x)$  for all  $x \in X$ , then  $O_f$  is the least element of  $\downarrow f$ . Now take  $g_1, g_2 \in \downarrow f$  and define the mapping  $h: X \rightarrow L$  by  $h(x) = g_1(x) \vee_{f(x)} g_2(x)$  for all  $x \in X$ , we claim  $h = g_1 \vee_f g_2$ .

Take an arbitrary  $a \in L$  and  $x \in h_{\langle\langle a \rangle\rangle}$ , then one can find a element  $\bar{a} \in L$  such that  $h(x) \gg \bar{a} \gg a$ . By Lemma 2.5(1) and  $h(x) = \bigvee_{f(x)} (\downarrow g_1(x) \cup \downarrow g_2(x))$ , there are finite subsets  $\{a_1, a_2, \dots, a_k\} \subseteq \downarrow g_1(x)$ ,  $\{b_1, b_2, \dots, b_m\} \subseteq \downarrow g_2(x)$  such that

$$h(x) \gg \bigvee_{f(x)} \{a_i, b_j: i = 1, 2, \dots, k; j = 1, 2, \dots, m\} = e \gg \bar{a}.$$

Hence  $f(x) \gg e$ . Let

$$V = \bigcap_{i=1}^k g_{1\langle\langle a_i \rangle\rangle} \cap \bigcap_{i=1}^k g_{2\langle\langle b_i \rangle\rangle} \cap f_{\langle\langle e \rangle\rangle},$$

then  $x \in V \in \Omega(X)$ . For each  $y \in V$ , as

$$h(y) = g_1(y) \vee_{f(y)} g_2(y) \gg \bigvee_{f(y)} \{a_i, b_j: i = 1, 2, \dots, k, j = 1, 2, \dots, m\}$$

and  $f(y) \gg e$ , by Lemma 2.5(2)

$$e = \bigvee_{f(y)} \{a_i, b_j: i = 1, 2, \dots, k, j = 1, 2, \dots, m\},$$

which implies  $y \in h_{\langle\langle a \rangle\rangle}$ , i.e.,  $V \subseteq h_{\langle\langle a \rangle\rangle}$ . By arbitrariness of  $x \in h_{\langle\langle a \rangle\rangle}$  we have  $h_{\langle\langle a \rangle\rangle} \in \Omega(X)$  and hence  $h \in [X \rightarrow L]$ .

Now suppose  $g \in \downarrow f$  such that  $g_1 \leq g$  and  $g_2 \leq g$ , then  $g(x) \geq g_1(x) \vee_{g(x)} g_2(x)$  for all  $x \in X$ . Again by Lemma 2.5(2) we have  $g(x) \geq g_1(x) \vee_{f(x)} g_2(x) = h(x)$  as  $f(x) \geq g(x)$ , so  $h = g_1 \vee_f g_2$  and the lemma is proved.

**Theorem 2.8.** Let  $L$  be a DCPO. Then  $L$  is a continuous  $L$ -Domain if and only if  $[X \rightarrow L]$  is a continuous  $L$ -Domain for all core compact spaces  $X$ .

**Proof.** The necessity of the theorem follows from Lemmas 2.6 and 2.7. However,  $L \cong [X \rightarrow L]$  if we take  $X$  as one-point space, so the sufficiency of the theorem follows.  $\square$

From the proof of Lemma 2.7, the following corollary is obvious.

**Corollary 2.9.** For a DCPO  $L$ ,  $L$  is bounded complete continuous if and only if  $[X \rightarrow L]$  is bounded complete continuous for all core compact spaces  $X$ .

In order to prove Theorem A, we need some notions. Following [3], we say that a DCPO  $L$  has property  $m$  if for each nonempty finite set  $A \subseteq L$  and each upper bound  $x_0$

of  $A$ , there is a minimal upper bound  $x$  of  $A$  such that  $x \leq x_0$ . A DCPO  $L$  is bicomplete if each filtered set in  $L$  has an infimum.

**Lemma 2.10.** *Each bicomplete DCPO  $L$  has property  $m$ .*

**Proof.** Let  $A$  be a nonempty finite set of  $L$  and  $x_0$  an upper bound of  $A$ , and write  $\mu(A) = \{x \in L : x \text{ is an upper bound of } A\}$  and  $G(A) = \{B \subseteq \mu(A) : B \text{ is a filtered set}\}$ ;  $G(A)$  is clearly a nonempty poset satisfying the condition of Zorn's Lemma as  $\{x_0\} \in G(A)$ . Hence  $G(A)$  has a maximal element  $B_0$  and  $B_0$  has the infimum  $b_0$  which is obviously a minimal upper bound of  $A$  with  $b_0 \leq x_0$ .  $\square$

The following lemma appears as “Theorem 1.37” in [3], we omit the proof.

**Lemma 2.11** [3]. *A DCPO  $L$  is bicomplete if  $[L \rightarrow L]$  is continuous.*

For DCPOs  $L$  and  $L_1$ , a continuous mapping  $r : L \rightarrow L_1$  is called a retraction if there is a continuous mapping  $h : L_1 \rightarrow L$  such that  $r \circ h$  equals the identity mapping on  $L_1$ . In this case  $L_1$  is called a retract of  $L$  and  $(r, h)$  a retraction embedding pair between  $L$  and  $L_1$ .

**Lemma 2.12.** *Any retract of an L-Domain (continuous DCPO) is again an L-Domain (continuous DCPO), and  $[X \rightarrow L_1]$  is a retract of  $[X \rightarrow L]$  if  $L_1$  is a retract of  $L$  for all topological spaces  $X$ .*

**Proof.** Suppose that  $L_1$  is a retract of  $L$  and  $(r, h)$  a retraction embedding pair between  $L$  and  $L_1$ . Note that both  $r$  and  $h$  are order-preserving and  $\downarrow x = \downarrow r \circ h(x) = r(\downarrow h(x))$  for each  $x \in L_1$ ; the completeness of  $\downarrow x$  follows from the completeness of  $\downarrow h(x)$ . Hence  $L_1$  is an L-Domain if  $L$  is. Also note that for each  $x \in L_1$ ,  $r(\downarrow h(x)) = \{r(y) : y \in \downarrow h(x)\} \subseteq \downarrow x$ ; then  $\downarrow x$  is directed and  $\bigvee \downarrow x = x$  if  $L$  is continuous. Now for each  $f \in [X \rightarrow L]$  and  $g \in [X \rightarrow L_1]$  define mappings  $\bar{r} : [X \rightarrow L] \rightarrow [X \rightarrow L_1]$  and  $\bar{h} : [X \rightarrow L_1] \rightarrow [X \rightarrow L]$  by  $\bar{r}(f) = r \circ f$  and  $\bar{h}(g) = h \circ g$ ; it is easy to verify that  $(\bar{r}, \bar{h})$  is a retraction embedding pair between  $[X \rightarrow L]$  and  $[X \rightarrow L_1]$ . Thus the lemma is proved.  $\square$

In [3] Jung gave a characterization of continuous L-Domains with least element in terms of retracts; we now give similarly a characterization of continuous L-Domains without least element.

**Lemma 2.13.** *Let  $L$  be a continuous DCPO with continuous mapping space  $[L \rightarrow L]$ . Then  $L$  is not an L-Domain if and only if  $L$  contains  $T_1$  or  $M_1$  as a retract, where  $T_1$  and  $M_1$  are continuous DCPOs denoted by Fig. 2 and Fig. 3 respectively.*

**Proof.** It is obvious that an L-Domain can't contain  $T_1$  or  $M_1$  as a retract because neither  $T_1$  nor  $M_1$  is an L-Domain, hence the sufficiency of the lemma holds. Now suppose that  $L$  is not an L-Domain. Then there is an  $x \in L$  such that  $\downarrow x$  has no least element or  $\downarrow x$  contains a pair of elements  $\{x_1, x_2\}$  without supremum in  $\downarrow x$ . We claim:




Fig. 2.  $T_1$




Fig. 2'.




Fig. 3.  $M_1$

(1) If  $\downarrow x$  has no least element, then  $L$  contains  $T_1$  as a retract.



By Lemma 2.11  $\downarrow x$  has at least two minimal elements  $\{\bar{x}_1, \bar{x}_2\}$  which are also minimal elements of  $L$ . Also by Lemma 2.10 and Lemma 2.11  $\{\bar{x}_1, \bar{x}_2\}$  has a minimum upper bound  $x_0 \leq x$ . Define a mapping  $r: L \rightarrow \{x_0, \bar{x}_1, \bar{x}_2\}$  by

$$r(x) = \begin{cases} x_0, & x \notin \{\bar{x}_1, \bar{x}_2\}, \\ x, & x \in \{\bar{x}_1, \bar{x}_2\}; \end{cases}$$

it is not hard to see that  $r$  is a retraction and its image  $r(L) = \{x_0, \bar{x}_1, \bar{x}_2\}$  is a copy of  $T_1$  inside  $L$ , so  $L$  contains  $T_1$  as a retract.

(2) If  $\{x_1, x_2\}$  has no supremum in  $\downarrow x$ , then  $L$  contains  $M_1$  as a retract.

By (1) we can assume that  $\downarrow x$  has the least element  $O_x$ , thus (2) follows from Jung's characterization [3, Proposition 4.22] and hence the lemma holds.  $\square$

Fig. 4.  $T_N$ Fig. 5.  $M_N$ 

**Lemma 2.14.** *Neither  $[T_N \rightarrow T_1]$  nor  $[M_N \rightarrow M_1]$  is continuous, where  $T_N$  and  $M_N$  are continuous DCPOs denoted by Fig. 4 and Fig. 5 respectively.*

**Proof.** Let  $f_{x_1} \in [T_N \rightarrow T_1]$  be the constant mapping taking value  $x_1$ . It is not hard to see  $f_{x_1} \notin \Downarrow f_{x_1}$ ; hence  $\Downarrow f_{x_1} = \emptyset$  because  $f_{x_1}$  is a minimal element of  $[T_N \rightarrow T_1]$ . Thus  $[T_N \rightarrow T_1]$  is not continuous.

In order to show that  $[M_N \rightarrow M_1]$  is not continuous, define mappings  $f, g_1, g_2 : M_N \rightarrow M_1$  as follows:

$f(O_N) = O_{M_1}$ ,  $f(y_1) = a_1$ ,  $f(y_2) = a_2$  and  $f(C_n) = 1_{M_1}$  for all  $n \in \mathbb{N}$ ;

$g_1(O_N) = O_{M_1}$ ,  $g_1(y_1) = a_1$ ,  $g_1(y_2) = O_{M_1}$  and  $g_1(C_n) = a_1$  for all  $n \in \mathbb{N}$ ;

$g_2(O_N) = O_{M_1}$ ,  $g_2(y_1) = O_{M_1}$ ,  $g_2(y_2) = a_2$  and  $g_2(C_n) = a_2$  for all  $n \in \mathbb{N}$ .

Obviously  $f, g_1, g_2 : [M_N \rightarrow M_1]$ . We can easily show  $f \notin \Downarrow f$  and  $g_1, g_2 \in \Downarrow f$ . Now we claim that  $\{g_1, g_2\}$  has no upper bound in  $\Downarrow f$ , i.e.,  $\Downarrow f$  is not directed and hence  $[M_N \rightarrow M_1]$  is not continuous.

Let  $h_1, h_2 \in [M_N \rightarrow M_1]$  be the mappings taking values  $\bar{a}_1$  and  $\bar{a}_2$  on the set  $\{c_n : n \in \mathbb{N}\}$  respectively,  $h_1(y_1) = h_2(y_1) = a_1$  and  $h_1(y_2) = h_2(y_2) = a_2$ . Obviously  $h_i \notin \Downarrow f$  ( $i = 1, 2$ ). Note that if  $g \leq f$  and  $g$  is an upper bound of  $\{g_1, g_2\}$ , then  $g \geq h_1$  or  $g \geq h_2$  and hence  $g \notin \Downarrow f$ . Thus the claim holds and hence the lemma is proved.  $\square$

**Theorem 2.15** (Theorem A). *For DCPO  $L$ , the following conclusions are equivalent:*

- (1)  $L$  is a continuous  $L$ -Domain;
- (2)  $[X \rightarrow L]$  is a continuous  $L$ -Domain for all core compact spaces  $X$ ;
- (3)  $[X \rightarrow L]$  is a continuous DCPO for all core compact spaces  $X$ .



**Proof.** (1)  $\Rightarrow$  (2) follows from Theorem 2.8, (2)  $\Rightarrow$  (3) is obvious, and we now prove (3)  $\Rightarrow$  (1).

In fact  $L$  is a continuous DCPO because  $L \cong [X \rightarrow L]$  for the one-point space  $X$ . Now suppose that  $L$  is not an L-Domain. By Lemma 2.13  $L$  contains  $T_1$  or  $M_1$  as a retract, and then by Lemma 2.12  $[X \rightarrow T_1]$  or  $[X \rightarrow M_1]$  is a retract of  $[X \rightarrow L]$  for all core compact spaces  $X$ . Especially, either  $[T_N \rightarrow T_1]$  is a retract of  $[T_N \rightarrow L]$  or  $[M_N \rightarrow M_1]$  is a retract of  $[M_N \rightarrow L]$ . Again by Lemma 2.12 either  $[T_N \rightarrow T_1]$  or  $[M_N \rightarrow M_1]$  is a continuous DCPO, which contradicts Lemma 2.14. Hence (3)  $\Rightarrow$  (1) holds.  $\square$

The following proposition is given by the referee.

**Proposition 2.16.**  $[X \rightarrow T_1]$  is a continuous DCPO for all spaces  $X$  which are both compact and core compact, where  $T_1$  is the continuous DCPO denoted by Fig. 2.

**Proof.** Let  $X$  be a core compact and compact space and consider the four-point lattice  $L$  (given by Fig. 2') with  $T_1$  consisting of the top three points. Then  $[X \rightarrow L]$  is a continuous lattice, hence a continuous DCPO. By the compactness of  $X$ , we can easily show that  $[X \rightarrow T_1]$  is a Scott open subset of  $[X \rightarrow L]$ . Hence  $[X \rightarrow T_1]$  is a continuous DCPO.  $\square$

Note that  $T_1$  is not an L-Domain; this proposition shows that the second part of Problem A has a negative answer.

### 3. Solution to Problem B

The Isbell topology on  $[X \rightarrow L]$ , written  $\text{Is}(X, L)$ , means the topology obtained by taking as a subbase for the open sets all sets of the form

$$N(H, V) = \{f \in [X \rightarrow L] : f^{-1}(V) \in H\},$$

where  $H$  is a Scott open set in  $\Omega(X)$  and  $V$  is a Scott open set in  $L$ .

**Example 3.1.** Let  $L$  be the DCPO denoted by Fig. 6.  $L$  is obviously a bounded complete continuous DCPO without least element. Take  $X = N$  equipped the discrete topology, then  $X$  is a core compact space. We claim that the Isbell and Scott topologies on  $[X \rightarrow L]$  do not agree.

In fact, by Theorem 2.8  $[X \rightarrow L]$  is a continuous L-Domain and hence  $\{\uparrow f : f \in [X \rightarrow L]\}$  is a base of the Scott topology on  $[X \rightarrow L]$ . Define mappings  $f, g : X \rightarrow L$  as follows:

$$g(1) = a_1, g(2) = a_2 \text{ and } g(n) = a_3 \text{ for all } n \geq 3;$$

$$f(1) = a_1, f(2) = O_2 \text{ and } f(n) = O_3 \text{ for all } n \geq 3,$$





Fig. 6.

Fig. 7.  $L(N)$ 

obviously  $f, g \in [X \rightarrow L]$  and  $g \in \uparrow f$ . Also note that

$$\{\uparrow F = \{B \subseteq N : B \supseteq F\} : F \text{ is a finite subset of } N\}$$

is a base of the Scott topology  $\sigma(\Omega(X))$ ; we can show that

$$B(g) = \left\{ N(\uparrow\{1\}, \{a_1\}) \cap N(\uparrow\{2\}, \{a_2\}) \cap N(\uparrow F, \{a_3\}) : \right. \\ \left. F \text{ is a finite subset of } N \setminus \{1, 2\} \right\}$$

is an Isbell neighbourhood base of  $g$ . Take an arbitrary

$$V = N(\uparrow\{1\}, \{a_1\}) \cap N(\uparrow\{2\}, \{a_2\}) \cap N(\uparrow F, \{a_3\}) \in B(g)$$

and let  $j_0 = \max F$ , define a mapping  $h : N \rightarrow L$  by  $h(1) = a_1$ ,  $h(2) = a_2$ ,  $h(j_0 + 1) = O_2$  and  $h(n) = a_3$  for all  $n \in \mathbb{N} \setminus \{1, 2, j_0 + 1\}$ . It is easy to see  $h \in V$  but  $h \notin \uparrow f$  as  $h(j_0 + 1) = O_2 \not\geq f(j_0 + 1) = O_3$ . So  $V \not\subseteq \uparrow f$  and hence by arbitrariness of  $V \uparrow f$  is not Isbell open, i.e.,  $\text{Is}(X, L) \neq \sigma([X \rightarrow L])$ .

**Example 3.2.** Let  $L(N)$  be the continuous L-Domain with least element  $O$  denoted by Fig. 7 and  $X = (L(N), \sigma(L(N)))$ . By Theorem 2.8  $[X \rightarrow L(N)]$  is a continuous L-Domain with least element.

Note that both  $L$  and  $\Omega(X)$  are countably based continuous DCPO, so  $\text{Is}(X, L)$  has a countable base. However  $\sigma([X \rightarrow L])$  doesn't have a countable base and hence  $\text{Is}(X, L) \neq \sigma([X \rightarrow L])$ .

From the definition of bounded complete DCPO, the following lemma is clear.

**Lemma 3.3.** A DCPO  $L$  is bounded complete if and only if each pair of elements with upper bound has a supremum.

**Lemma 3.4.** *Let  $L$  be a continuous  $L$ -Domain with least element  $O_L$ . Then  $L$  is a bounded complete DCPO if  $\text{Is}(X, L) = \sigma([X \rightarrow L])$  for all core compact spaces  $X$ .*

**Proof.** Suppose that  $L$  is not bounded complete. By Lemma 3.3 there is a pair of elements  $\{b_1, b_2\}$  with upper bound which has no supremum, which implies  $\{b_1, b_2\}$  has at least two minimal upper bounds  $\{a_1, a_2\}$  and  $\uparrow a_1 \cap \uparrow a_2 = \emptyset$  because  $L$  is an  $L$ -Domain. By Theorem 1.1 and continuity of  $L$  we can find the elements  $b'_1$  and  $b'_2$  with  $b'_1 \ll b_1$  and  $b'_2 \ll b_2$  such that  $\{b'_1, b'_2\}$  has no supremum, then both  $a_1^* = b'_1 \bigvee_{a_1} b'_2$  and  $a_2^* = b'_1 \bigvee_{a_2} b'_2$  are minimal upper bounds of  $\{b'_1, b'_2\}$  and  $\uparrow a_1^* \cap \uparrow a_2^* = \emptyset$ . Now take  $X = L(N)$  (denoted by Fig. 7) equipped with the Scott topology, and define mappings  $f, g: X \rightarrow L$  as follows:

$$f(O) = O_L, \quad f(z_1) = b'_1, \quad f(z_2) = b'_2 \quad \text{and} \quad f(d_n) = \begin{cases} a_1^*, & n \text{ an odd number,} \\ a_2^*, & n \text{ an even number,} \end{cases}$$

$$g(O) = O_L, \quad g(z_1) = b_1, \quad g(z_2) = b_2 \quad \text{and} \quad g(d_n) = \begin{cases} a_1, & n \text{ an odd number,} \\ a_2, & n \text{ an even number,} \end{cases}$$

we can verify that  $g, f \in [X \rightarrow L]$  and  $f \ll g$ . By Theorem 2.8  $\uparrow f$  is a Scott open neighbourhood of  $g$ . Write  $\mathcal{F}_1$  and  $\mathcal{F}_2$  for the sets of all nonempty finite subsets of  $\{x_{2n-1}: n \in \mathbb{N}\}$  and  $\{x_{2n}: n \in \mathbb{N}\}$  respectively, and let

$$B(g) = \left\{ N(\uparrow F_1, \uparrow t_1) \cap N(\uparrow F_2, \uparrow t_2) \cap N(\uparrow z_1, \uparrow t_3) \cap N(\uparrow z_2, \uparrow t_4) : \right. \\ \left. F_1 \in \mathcal{F}_1, F_2 \in \mathcal{F}_2 \text{ and } a_1^* \leq t_1 \ll a_1, \quad a_2^* \leq t_2 \ll a_2, \right. \\ \left. t_3 \in \downarrow b_1 \setminus \downarrow b_2, \quad t_4 \in \downarrow b_2 \setminus \downarrow b_1 \right\}.$$

Then  $B(g)$  is an Isbell neighbourhood base of  $g$ . Take an arbitrary

$$V = N(\uparrow F_1, \uparrow t_1) \cap N(\uparrow F_2, \uparrow t_2) \cap N(\uparrow z_1, \uparrow t_3) \cap N(\uparrow z_2, \uparrow t_4) \in B(g)$$

and let  $j_0 = \max\{i \in \mathbb{N}: x_i \in F_1 \cup F_2\}$ , and define mapping  $h: X \rightarrow L$  as follows:

$$h(x) = g(x) \quad \text{for all } x \in L(N) \setminus \{d_{2j_0+1}\} \quad \text{and} \quad h(d_{2j_0+1}) = a_2.$$

It is not hard to see  $h \in [X \rightarrow L]$  and  $h \in V$ , but  $h \notin \uparrow f$  as  $h(x_{2j_0+1}) = a_2 \not\leq a_1^*$  (otherwise  $a_2 \in \uparrow a_1^* \cap \uparrow a_1^* = \emptyset$ ). Hence  $V \not\subseteq \uparrow f$ , which implies that  $\uparrow f$  is not an Isbell open set, so  $\text{Is}(X, L) \neq \sigma([X \rightarrow L])$ . This is a contradiction and hence the lemma is proved.  $\square$

**Theorem 3.5** (Theorem B). *Let  $L$  be a continuous  $L$ -Domain with least element  $O_L$ . Then  $\text{Is}(X, L) = \sigma([X \rightarrow L])$  for all core compact spaces  $X$  if and only if  $L$  is a bounded complete DCPO.*

**Proof.** The necessity of the theorem follows from Lemma 3.4. Conversely, note that  $\text{Is}(X, L) \subseteq \sigma([X \rightarrow L])$ . By Theorem 1.1 we need only show that  $\uparrow f$  is an Isbell open set for each  $f \in [X \rightarrow L]$ . Write  $bV$  for the mapping taking value  $b$  on  $V$  and value  $O_L$  otherwise, and take an arbitrary  $f \in [X \rightarrow L]$ .

For each  $g \in \uparrow f$ , by Lemma 2.6 we have

$$g = \bigvee_{a \in L} \bigvee_{b \in \downarrow a} \bigvee_{V \in \downarrow g_{\langle\langle a \rangle\rangle}} F(b, V, g) = \bigvee_{a \in L} \bigvee_{b \in \downarrow a} \bigvee_{V \in \downarrow g_{\langle\langle a \rangle\rangle}} bV.$$

Then by Corollary 2.9 there exists a finite subset  $\{b_1, b_2, \dots, b_k\} \subseteq L$  such that  $f \leq \bigvee \{b_i V_i : i = 1, 2, 3, \dots, k\}$ , where  $b_i \in \downarrow a_i$  and  $V_i \in \downarrow g_{\langle\langle a_i \rangle\rangle}$  ( $\forall i = 1, 2, \dots, k$ ). Obviously  $H = \bigcap_{i=1}^k N(\uparrow V_i, \uparrow a_i)$  is an Isbell neighbourhood of  $g$ , we claim  $H \subseteq \uparrow f$ :

In fact, for each  $h \in H$ , by Corollary 2.9 and Lemma 2.4 we have

$$f \leq \bigvee \{b_i V_i : i = 1, 2, \dots, k\} \ll \bigvee \{a_i h_{\langle\langle a \rangle\rangle} : i = 1, 2, \dots, k\} \leq h$$

because  $V_i \ll h_{\langle\langle a_i \rangle\rangle}$  and  $b_i \ll a_i$  ( $i = 1, 2, \dots, k$ ), so  $h \in \uparrow f$ . By the arbitrariness of  $g \in \uparrow f$ , it follows that  $\uparrow f$  is Isbell open and thus the theorem is proved.  $\square$

By Examples 3.1 and 3.2 the condition that  $L$  is a continuous L-Domain with least element in Theorem 3.5 is suitable.

## Acknowledgement

The authors gratefully acknowledge the referee for his careful reading and offering helpful suggestions and corrections. Also we would like to thank Dr. A. Jung for sending us interesting papers.

## References

- [1] J. van Mill and G.M. Reed, eds., Open Problems in Topology (North-Holland, Amsterdam, 1990).
- [2] J.D. Lawson, The versatile continuous order, in: Mathematical Foundation of Programming Semantics, Lecture Notes in Computer Science 298 (Springer, Berlin, 1988) 134–160.
- [3] A. Jung, Cartesian closed categories of domains, Ph.D. Thesis, Technische Hochschule Darmstadt.
- [4] F. Schwarg and S. Weck, Scott topology, Isbell topology and continuous convergence, in: Continuous Lattices and Their Applications, Lecture Notes in Pure and Applied Mathematics 101 (Dekker, New York, 1985) 251–273.
- [5] G. Gierz et al., A Compendium of Continuous Lattices (Springer, Berlin, 1980).